

Hadrons Fall2023

HW1 due 27/10/2023

(+1 points for handing in on time)

Q1 Black-body radiation

- a) The energy spectrum of photons in the momentum space reads

$$u_E(k, T) \propto k^2 \frac{k}{e^{k/T} - 1}.$$

Show that the maximum of $u_E(k, T)$ occurs at

$$(3 - k/T) e^{k/T} - 3 = 0.$$

Solve the equation numerically and verify that $k/T \approx 2.8$. This is Wien's displacement Law.

- b) Repeat the above exercise but instead work in the wavelength space.

Show that

$$u_\lambda(\lambda, T) \propto \frac{1}{\lambda^5} \frac{1}{e^{\frac{2\pi}{\lambda T}} - 1}.$$

Derive the corresponding Wien's Law.

- c) The surface temperature of the sun is 5778 K. Find the “most likely” energies of solar radiation according to the two versions of Wien's Law. Why are they **NOT** the same?

Q2 Particle adventure.

Visit the PDG website [link](#) to look up particle properties.

- a) Construct a table of the masses of the lowest vector mesons (1 – – state): $(u\bar{u} = d\bar{d}, s\bar{s}, c\bar{c}, b\bar{b})$, alongside the current quark masses.
b) Look up the Fermi coupling constant G_F and mass of W-boson M_W .
c) Given

$$\frac{G_F}{\sqrt{2}} = \frac{g_W^2}{8M_W^2},$$

what is the value of the weak interaction coupling constant g_W and $\alpha_W = \frac{g_W^2}{4\pi}$? Compare this to the QED value

$$\alpha_{EM} = \frac{1}{137},$$

which one is stronger?

Q3 Kinematics.

- a) Consider the scattering of two particles (of mass m_1 and m_2) in the Center-of-Mass (COM) frame. The energy in this frame is also the invariant mass \sqrt{s} of the system, i.e.

$$E_{\text{COM}} = \sqrt{s} = \sqrt{q^2 + m_1^2} + \sqrt{q^2 + m_2^2}.$$

Show that

$$q(s) = \frac{1}{2}\sqrt{s}\sqrt{1 - \frac{(m_1 + m_2)^2}{s}}\sqrt{1 - \frac{(m_1 - m_2)^2}{s}}.$$

- b) Show that the invariant mass (\sqrt{s}) of a 3-body system satisfies

$$s = s_{12} + s_{23} + s_{13} - m_1^2 - m_2^2 - m_3^2,$$

where standard relativistic kinematics apply, i.e.

$$\begin{aligned} s_{ij} &= (p_i + p_j)^2 \\ p_i &= (E_i, \vec{p}_i) \\ p_i^2 &= E_i^2 - (\vec{p}_i)^2. \end{aligned}$$

Q4 N-body phase space.

The N-body Lorentz Invariant phase space (LISP) is defined as

$$\phi_N(s = P^2) = \int \frac{d^3p_1}{(2\pi)^3} \frac{1}{2E_1} \frac{d^3p_2}{(2\pi)^3} \frac{1}{2E_2} \cdots \frac{d^3p_N}{(2\pi)^3} \frac{1}{2E_N} \times (2\pi)^4 \delta^4(P - \sum_i p_i).$$

Note that $E_j = \sqrt{p_j^2 + m_j^2}$.

- a) Derive an analytic expression of the 2-body phase space. (in terms of s, m_1, m_2)
b) For all $m_i = 0$, derive an analytic expression of the 3-body phase space.

ans:

$$\phi_3(s) = \frac{s}{256\pi^3}.$$

c) The decay width of the muon can be estimated by

$$\Gamma = \frac{G_F^2 m_\mu^5}{192\pi^3}.$$

What is the interaction involved? Draw the Feynman diagram. Plug in numbers and calculate its lifetime. Do you understand the factor of m_μ^5 ?